

	M05.1 - EPINEPHRINE FOR ANAPHYLAXIS (1 mg/ml)	
	MEDICATION STANDING ORDER	
Version date: 2023-12-14	Effective date: 2024-01-16 (0700)	

INDICATIONS
<ul style="list-style-type: none"> Known or suspected anaphylaxis

CONTRAINDICATIONS
<ul style="list-style-type: none"> Not applicable

DOSING		
AGE	INITIAL DOSE	REPEAT DOSE
AUTOINJECTOR		EMR
6 yrs & older	<ul style="list-style-type: none"> 0.3 mg (orange) 	<ul style="list-style-type: none"> Once as required
12 mos up to 6 yrs	<ul style="list-style-type: none"> 0.15 mg 2 (green) 	
INTRAMUSCULAR		PCP / ICP
17 yrs & older	<ul style="list-style-type: none"> 0.5 mg 	<ul style="list-style-type: none"> Every 5 to 15 minutes as required
12 mos up to 17 yrs	<ul style="list-style-type: none"> 0.01 mg/kg (MAX = 0.5 mg) 	

NOTES
<ol style="list-style-type: none"> Confirm correct concentration & dose (QRG). If a pediatric-dose autoinjector is not available, use the adult device. Administer into the lateral thigh muscle (vastus lateralis). It has greater blood flow than the shoulder muscle (deltoid), resulting in faster absorption and higher drug levels.

APPROVED BY	
	
EMS Medical Director	EMS Associate Medical Director

VERSION CHANGES (refer to X08 for change tracking)
<ul style="list-style-type: none"> High-alert warning removed

QRG: INTRAMUSCULAR DOSING		
WEIGHT (kg)	DOSE (mg)	VOLUME (ml)
5 to 10	0.1	0.1
11 to 15	0.15	0.15
16 to 20	0.2	0.2
21 to 25	0.25	0.25
26 to 30	0.3	0.3
31 to 35	0.35	0.35
36 to 40	0.4	0.4
41 to 45	0.45	0.45
> 45	0.5	0.5