	M06.4 - GLUCAGON POWDER (<i>BAQSIMI</i>)	
	Version date: 2025-02-04	Effective date: 2025-04-30 (07:00)



INDICATIONS
<ul style="list-style-type: none"> Known or suspected hypoglycemia ¹

WARNINGS
ABSOLUTE CONTRAINDICATIONS: <ul style="list-style-type: none"> Hypersensitivity to glucagon


ADMINISTRATION	
INTRANASAL	EMR / PCP / ICP / ACP
4 YEARS & OLDER: <ul style="list-style-type: none">• 3 mg once	

NOTES
<ol style="list-style-type: none"> If blood glucose testing is not available, treat based on symptoms. If there is no response within 15 minutes, glucose or dextrose must be administered. Glucagon may cause significant nausea. Consider co-administration of antiemetic. Ensure that the patient is eating or receives oral glucose / intravenous dextrose after glucagon administration.

LINKS
<ul style="list-style-type: none"> E10 - Hypoglycemia

APPROVED BY	
	
Provincial Medical Director	Provincial Associate Medical Director

VERSION CHANGES (refer to X08 for change tracking)
<ul style="list-style-type: none"> Addition of advanced work scope

APPENDIX A: ADMINISTRATION OF <i>BAQSIMI</i> GLUCAGON NASAL POWDER.	
STEP #1: Remove the shrink wrap by pulling on the red stripe. Open the lid & remove the device from the tube. DO NOT TEST BEFORE USE.	
STEP #2: Hold the device between your thumb and fingers. DO NOT PRESS THE PLUNGER UNTIL YOU ARE READY TO ADMINISTER.	
<p>STEP #3: Insert the tip gently into one nostril until your fingers touch the outside of the nose.</p> 	<p>STEP #4: Push the plunger all the way in. The dose is complete when the green line is no longer showing.</p> 