

	INDICATIONS	
HyperkalemiaMagnesium toxicity		

CONTRAINDICATIONS	
• None	

MAGNESIUM TOXICITY				
ROUTE (WORK SCOPE)	INITIAL DOSE	REPEAT DOSE		
INTRAVENOUS (PCP & ABOVE)		Neze		
INTRAOSSEOUS (ICP & ABOVE)	10 years & older - 1 gram (10 ml)	None		

HYPERKALEMIA		
ROUTE (WORK SCOPE)	INITIAL DOSE	REPEAT DOSE
INTRAVENOUS / INTRAOSSEOUS (ICP & ABOVE)	10 years & older - 1 to 2 gm (10 - 20 ml)	Once in 5 minutes if ECG signs persist & every 30 - 60 minutes if ECG signs recur

NOTES	
1.	ERS HIGH-ALERT MEDICATION: Refer to Shared Health Provincial Clinical Standard - Safety Controls for High-Alert Medications (refer to A03 - HIGH ALERT MEDICATIONS).
2.	During cardiac arrest administer by rapid push followed by saline flush.
3.	For magnesium toxicity administer by slow push over 3 to 5 minutes

APPROVED BY	
Buftsterel	Jefferman L.
Medical Director - Provincial EMS/PT	Associate Medical Director – Provincial EMS/SPT

VERSION CHANGES (refer to X08 for change tracking)

• Addition of Shared Health Provincial Clinical Standard for high-alert medications